| For office use only | | | | | |
|---------------------|--------------|--|--|--|--|
| Date Received | Date Replied | | | | |
| | | | | | |

MGA NITROGEN PREDICTOR Data Sheet

To calculate the nitrogen requirement for your individual maize field/s we need to know details of present and past field management. Please complete your name, farm and contact details in the spaces provided, then input your field details *in one box only*.

Once complete please send this form to the MGA Office, Town Barton Farm, Sandford, Crediton, Devon. EX17 4LS. Or Fax: 01363 776938. Should you have any queries please contact the office. Tel: 01363 775040.

| L | | | l l | | | <u> </u> | | | | |
|---|--|---------|---------|---------|---------|----------|---------|---------|--|--|
| | | E: 114 | F: 110 | F: 110 | I =: | I =: | F: 110 | E: | | |
| | | Field 1 | Field 2 | Field 3 | Field 4 | Field 5 | Field 6 | Field 7 | | |
| Field Details | Field Name | | | | | | | | | |
| | Field Size (acres) | | | | | | | | | |
| | Variety | | | | | | | | | |
| | Typical Nitrogen application u/acre | | | | | | | | | |
| | 31 5 11 | | | | | | | | | |
| Variety | Late Maturing | | | | | | | | | |
| | Mid Maturing | | | | | | | | | |
| | • | | | | | | | | | |
| | Early Maturing | | | | | | | | | |
| | | | | | | | | | | |
| Harvest Date | Plus 2 weeks early | | | | | | | | | |
| | Normal | | | | | | | | | |
| | Plus 2 weeks late | | | | | | | | | |
| | | | | | | | | | | |
| Soil Type | Heavy | | | | | | | | | |
| ,, | Medium | | | | | | | | | |
| | Light | | | | | | | | | |
| | Ligiti | | | | | | | | | |
| Coil Ctructure | (Cuba silad this user) Cood | | | | | | | | | |
| Soil Structure | (Subsoiled this year) Good | | | | | | | | | |
| | (subsoiled in last 2 or 3 years) Average | | | | | | | | | |
| | (not subsoiled recently) Poor | | | | | | | | | |
| | | | | | | | | | | |
| Previous Cropping | Maize | | | | | | | | | |
| | Cereals | | | | | | | | | |
| | Grass (1-2 years) | | | | | | | | | |
| | 3 years Grass | | | | | | | | | |
| | 3 + years Grass | | | | | | | | | |
| | Rape/Beans/Other break | | | | | | | | | |
| | Rape/Beans/Other break | | | | | | | | | |
| 5 | | | | | | | | | | |
| Past history for | Arable rotation - no organic | | | | | | | | | |
| Last 10 years | manure | | | | | | | | | |
| | Arable rotation + organic manure | | | | | | | | | |
| (Continuous maize should be classified | Stock/Maize/Grass Rotation | | | | | | | | | |
| as arable rotation) | Stock/Maize/Grass rotation plus | | | | | | | | | |
| as arabic rotation) | organic manure | | | | | | | | | |
| | | | | | | | | | | |
| Spring & Autumn | High rainfall | | | | | | | | | |
| weather before drilling | (*1.5 + normal) average | | | | | | | | | |
| | Average | | | | | | | | | |
| | Low Rainfall | | | | | | | | | |
| | (*1.5 - normal) average | | | | | | | | | |
| | | | | | | | | | | |
| Drilling Date | Early April | | | | | | | | | |
| | 14/4 - 14/5 | | | | | | | | | |
| | 14/5 + Later | | | | | | | | | |
| | , c . Later | | l | l | l | l | L | | | |

| FYM | | Field 1 | Field 2 | Field 3 | Field 4 | Field 5 | Field 6 | Field 7 |
|---------------------------|----------------------|---------|---------|----------|---------|---------|---------|---------|
| Cattle | Tons/acre | | | | | | | |
| Applied before Feb | | | | | | | | |
| | Tonnes/ha | | | | | | | |
| Applied offer Feb | Tana/aara | | | | | | | |
| Applied after Feb | Tons/acre | | | | | | | |
| | Tonnes/ha | | | | | | | |
| | | | | | | | | |
| Pig | | | | | | | | |
| Applied before Feb | Tons/acre | | | | | | | |
| | Tonnes/ha | | | | | | | |
| | 1011163/114 | | | | | | | |
| Applied after Feb | Tons/acre | | | | | | | |
| | T // | | | | | | | |
| | Tonnes/ha | | | | | | | |
| Poultry (Layer manure) | | | | | | | | |
| (Laryer manus) | Tons/acre | | | | | | | |
| Applied before Feb | | | | | | | | |
| | Tonnes/ha | | | | | | | |
| Applied after Feb | Tons/acre | | 1 | + | | | - | |
| Applied diter I eb | 10115/4016 | | | | | | | |
| | Tonnes/ha | | | | | | | |
| 0 | | | | | | | | |
| Sewage Applied before Feb | النوال | | | | | | | |
| Applied before Feb | High | | | | | | | |
| | Low | | | | | | | |
| | | | | | | | | |
| Applied after Feb | High | | | | | | | |
| | Low | | | | | | | |
| | LOW | | | | | | | |
| Slurry | | | | | | | | |
| Cattle | Gallons/acre | | | | | | | |
| Applied before Feb | | | | | | | | |
| | M³/ha | | | | | | | |
| Applied after Feb | Gallons/acre | | | | | | | |
| | | | | | | | | |
| | M³/ha | | | | | | | |
| Pig | | | | | | | | |
| Applied before Feb | Gallons/acre | | | | | | | |
| / Applied Solote 1 os | Canono, acro | | | | | | | |
| | M³/ha | | | 1 | | | | |
| A . I' . I . G | 0 " ' | | 1 | 1 | | | | |
| Applied after Feb | Gallons/acre | | | | | | | |
| | M³/ha | | | + | | | | |
| | | | | <u> </u> | | | | |
| Poultry (broiler litter) | | | | | | | | |
| A - 15 - 11 - 6 - 5 - 5 | Tons/acre | | | | | | | |
| Applied before Feb | Tonnes/ha | | 1 | 1 | | | | |
| | 1011165/118 | | | | | | | |
| Applied after Feb | Tons/acre | | 1 | 1 | | | | |
| | | | 1 | 1 | | | | |
| | Tonnes/ha | | | | | | | |
| Cowere limit | | | | | | | | |
| Sewage liquid | Gallons/acre | | | | | | | |
| Applied before Feb | Gallolis/acte | | | | | | | |
| Applied befole Leb | M³/ha | | 1 | 1 | | | | |
| | | | | | | | | |
| Applied after Feb | Gallons/acre | | | | | | | |
| | M³/ha | | | 1 | | | | |
| | ivi ⁻ /na | | | | | | | |
| | | 1 | 1 | 1 | 1 | I | 1 | l . |